



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

289. Proposed by G. W. DROKE, Professor of Mathematics, University of Arkansas.

Find the curve such that the rectangle under the perpendiculars from two fixed points on the normals be constant.

Solution by S. G. BARTON, Ph. D., Clarkson School of Technology, Potsdam, N. Y.

This is problem 27, page 61, Murray's *Differential Equations*.

Take the straight line through the fixed points as the x axis, and the middle point as the origin. The coordinates of the points are then $(a, 0)$ and $(-a, 0)$. Using p for $\frac{dy}{dx}$, the equation of a normal at (x', y') is

$$y - y' = -\frac{1}{p}(x - x').$$

Its distance from $(a, 0)$ is

$$\frac{\frac{a}{p} - \left(y' + \frac{x'}{p}\right)}{\sqrt{1 + \frac{1}{p^2}}},$$

and its distance from $(-a, 0)$ is

$$\frac{-\frac{a}{p} - \left(y' + \frac{x'}{p}\right)}{\sqrt{1 + \frac{1}{p^2}}}.$$

The constant product, dropping primes, is

$$\frac{y^2 + \frac{2xy}{p} + \frac{x^2}{p^2} - \frac{a^2}{p^2}}{1 + \frac{1}{p^2}} = k^2,$$

$$\text{or } p^2 y^2 + 2xyp + x^2 - a^2 = k^2 p^2 + k^2, \quad \text{or } (y^2 - k^2)p^2 + 2xyp + x^2 - a^2 - k^2 = 0;$$

whence the p discriminant is

$$\text{Hence, } 4x^2 y^2 - 4(y^2 - k^2)(x^2 - a^2 - k^2).$$

$$4x^2 y^2 - 4x^2 y^2 + 4(k^2 + a^2)y^2 - k^2(x^2 - a^2 - k^2) = 0,$$

$$\text{or, } \frac{y^2}{k^2} + \frac{x^2}{k^2 + a^2} = 1.$$

This is the singular solution of the differential equation. It represents a system of confocal conics having the fixed points as foci.

Also solved by V. M. Spunar, G. B. M. Zerr, and J. Scheffer.

Solution by PROFESSOR F. L. GRIFFIN, Williams College.

However the ellipse is placed relative to the co-ordinate axes, its area is

$$S = \int_{x_1}^{x_2} (y'' - y') dx,$$

where x_1 and x_2 [$x_1 < x_2$] are the extreme abscissas taken in the curve, and y' and y'' [$y' \leq y''$] are the two ordinates corresponding to any one value of x . Solved for y , the given equation becomes

$$by = -(hx + f) \pm \sqrt{[f^2 - bc + 2(hf - bg)x + (h^2 - ab)x^2]}.$$

Now let

$$f^2 - bc = b^2 A, \quad fh - bg = b^2 B, \quad ab - h^2 = b^2 C,$$

where for an ellipse $C > 0$. Then the difference of the two ordinates becomes

$$y'' - y' = 2\sqrt{[A + 2Bx - Cx^2]}.$$

Hence, integrating,

$$S = 2 \left[\frac{Cx - B}{2C} \sqrt{[A + 2Bx - Cx^2]} + \frac{B^2 + AC}{2C^{\frac{3}{2}}} \sin^{-1} \frac{Cx - B}{\sqrt{[B^2 + AC]}} \right]_{x_1}^{x_2}$$

Now the extreme abscissas make $y' = y''$, or $A + 2Bx - Cx^2 = 0$; whence

$$Cx_2 = B + \sqrt{[B^2 + AC]} \quad \text{and} \quad Cx_1 = B - \sqrt{[B^2 + AC]}.$$

Substituting these values,

$$S = \frac{B^2 + AC}{C^{\frac{3}{2}}} [\sin^{-1} 1 - \sin^{-1}(-1)] = \frac{\pi}{b} \frac{[(fh - bg)^2 + (ab - h^2)(f^2 - bc)]}{(ab - h^2)^{\frac{3}{2}}},$$

which immediately reduces to the formula proposed.

Also solved by G. B. M. Zerr and J. Scheffer.

MECHANICS.

240. Proposed by S. A. COREY, Hiteman, Iowa.

A perfectly flexible wire rope weighing one pound per foot is suspended from the tops of two vertical supports 300 feet apart, one support being 30 feet higher than the other. One end of the rope is fastened to the top of the higher support, while 600 feet of the rope hangs vertically from the top of the lower support. Assuming that the rope is free to slide over the top of the lower support without friction, find the lowest point of